

Estreptoganabencil
Reg. Key S.A.G.A.R.P.A. Q-0524-014*

FOR VETERINARY USE

Penicillin - Streptomycin

Injectable broad-spectrum antibiotic powder and diluent reconstitution
with rapid and long-lasting effect.

DRUG DESCRIPTION

Each 1.5 mL of reconstituted product contains:

Penicillin G. Benzathine	150 000 I.U.
Penicillin G. Procaine	112 500 I.U.
Penicillin G. Sodium	37 500 I.U.
Streptomycin Sulfate	
Equivalent to Streptomycin Base	0.225 g
Monohydrated Sodium Dipyrone	220 mg
Guaiacol	11 mg
Excipient c.b.p.	1.5 ml

Estreptoganabencil® is an injectable broad-spectrum bactericidal antibiotic reconstitution with long-lasting effect, antipruritic and expectorant effective against Gram positive and Gram negative bacteria, the penicillins contained in this product belong to the Beta-lactamases group, the streptomycin another of its chemically active elements is included as an aminoglycoside.

Estreptoganabencil® is recommended as treatment of infections such as: mastitis, metritis, pneumonia, rhinitis and bacterial arthritis in Dairy Cattle, sheep, goats, swine, horses, cats and dogs. For barnyard fowl is recommended as treatment of ulcerative enteritis, borreliosis, psittacosis, colibacillosis and avian cholera.

Estreptoganabencil® achieves an antibiotic rapid and long-lasting effect, it can be used again after 48 to 72 hours, less injections are needed per treatment.

Estreptoganabencil® has a synergistic effect among its antibiotics which shows more effectiveness. It may be used in pregnant females with no risk of miscarriage. DRUG USE For infectious diseases caused by germs sensitive to Penicillin and Streptomycin in Dairy Cattle, horses, swine, sheep, goats, dogs, cats and barnyard fowl.

DOSAGE

Barnyard Fowl: 0.10 mL or 0.25 mL/kg.

Cats: 0.25 to 0.5 mL.

Dogs: 0.5 to 3 mL.

Dairy Cattle and Horses: 1 mL per 20 kg.

Swine, Sheep and Goats: 1 to 2 mL per 20 kg.
Frequency: Every 2 to 5 days according to prescription.

ROUTE OF ADMINISTRATION

Intramuscular

DRUG WARNINGS

Do not use in animals sensitive to Penicillins or Streptomycin. Do not use in horses destined for human consumption. Skin rash, pruritus and in acute cases anaphylaxis can be shown in hypersensitive animals. In case of allergic reaction stop use and treat with antihistamines.

WITHDRAWAL PERIOD

Milk withdrawal: 4 days

Meat withdrawal: 30 days

Poultry withdrawal: 4 days

Egg withdrawal: 48 hours.

Keep this product in a cool and dry place protected from light at a temperature between 15° and 30 °C

Once reconstituted the product can be stored up to 15 days in refrigeration with a temperature between 2o and 7o C. For reconstitution use the whole content in the vial and the included diluent.

Keep out of reach of children and pets.

Consult a veterinarian Prescription Only Medicine – Veterinarian (POM-V category)

PRESENTATION

1 vial x 4 000 000 I.U.

1 vial x 6 000 000 I.U.

REGISTRATION KEYS

Costa Rica MAG MX7-05-05-032

El Salvador VET.95-11-1068

Honduras PF-4588

Nicaragua 2373

Dominican Rep. 1475

MADE IN MEXICO BY: RIVERFARMA, S.A. DE C.V. Prol. Calz. de La Viga No. 1949, Col. Prado Churubusco, Del. Coyoacán. C.P.04230, México, D.F. E-mail: riverfarma@riverfarma.com.mx Web: www.riverfarma.com.mx Tel. 6389-6687, 6389-6688, 6389-6689. Within Mexico Toll Free: 01-800-505-7777